



Original Article

Relationship of left atrial size to obstructive sleep apnea severity in end-stage renal disease

Owen D. Lyons^{a,b}, Christopher T. Chan^c, Rosilene M. Elias^{a,b,c}, T. Douglas Bradley^{a,b,c,*}^a Sleep Research Laboratory, University Health Network Toronto Rehabilitation Institute, Toronto, ON, Canada^b Centre for Sleep Medicine and Circadian Biology, University of Toronto, Toronto, ON, Canada^c Department of Medicine, University Health Network Toronto General Hospital, Toronto, ON, Canada

ARTICLE INFO

Article history:

Received 14 February 2014

Received in revised form 24 June 2014

Accepted 3 July 2014

Available online 8 July 2014

Keywords:

Cardiovascular

End-stage renal disease

Fluid shift

Left atrium

Men

Obstructive sleep apnea

ABSTRACT

Background: Increased left atrial (LA) size is linked to elevated mortality in end-stage renal disease (ESRD). In addition, the degree of overnight rostral fluid shift from the legs is associated with severity of obstructive sleep apnea (OSA). As rostral fluid shift might distend the left atrium and increase fluid accumulation in the neck, we postulated that LA size would be related to the degree of overnight rostral fluid shift and OSA severity in ESRD patients.

Methods: Patients with ESRD underwent echocardiography and polysomnography. Leg fluid volume (LFV) was measured by bioelectrical impedance before and after the overnight sleep study in a subset of 21 patients.

Results: Forty patients (22 men), with a mean apnea–hypopnea index (AHI) of 25.1 ± 23.4 /h of sleep, had echocardiography and polysomnography performed. In men, there was a correlation between the AHI and LA size indexed for body surface area ($r = 0.743$, $p < 0.001$) that was not observed in women. Strong relationships were seen, again in men only, between LA size indexed to body surface area and the overnight change in leg fluid volume (Δ LFV) ($r = -0.739$, $p = 0.02$) and between AHI and Δ LFV ($r = -0.863$, $p = 0.003$).

Conclusions: In ESRD patients, there are relationships between Δ LFV and both LA size and OSA severity. These findings suggest that the relationship between LA size and mortality in ESRD may be related to Δ LFV and severity of OSA.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Patients with end-stage renal disease (ESRD) have an annual mortality rate between 10% and 20%, mainly due to cardiovascular causes [1–3]. Increased left atrial (LA) size contributes to cardiovascular mortality in both the general population and in patients with ESRD [4,5]. In ESRD, the prevalence of obstructive sleep apnea (OSA), characterized by repetitive upper airway collapse during sleep, is at least 50% and is much higher than in the general population [6,7] with men having a higher prevalence than women [8]. In ESRD patients on peritoneal dialysis, the presence of OSA has been shown to be an independent predictor of cardiovascular morbidity and death [9]. It has also been shown that the severity of OSA is directly related to the degree of overnight fluid shift from the legs in ESRD [10]. Given that ESRD causes fluid retention, that OSA is very common in ESRD,

and that OSA increases cardiovascular risk [2,3,11], it is important to examine potential mechanisms through which fluid retention, nocturnal rostral fluid shift, and OSA may contribute to potentially reversible changes in cardiac structure in ESRD patients.

The left atrium plays a crucial role in maintaining optimal cardiac function. It modulates left ventricular (LV) filling by acting as a reservoir during ventricular systole, a conduit during early diastole, and as an active pump in late diastole [12]. Its thin-walled structure means that it is more distensible than the left ventricle in response to increases in pressure or volume of pulmonary venous inflow [13]. The most common causes of an enlarged left atrium are LV dysfunction or hypertrophy, mitral valve disease, and fluid overload [13,14]. The relationship with fluid overload is of particular relevance in ESRD, given that fluid overload is a hallmark of this condition and predicts increased mortality [15].

The high prevalence of OSA in ESRD and in heart failure [16], both pathological states characterized by fluid overload, has led to an interest in the role of fluid retention and fluid shift from the legs in the pathogenesis of OSA in these conditions. In ESRD, the increased prevalence of OSA is not explained solely by comorbidities or increased body mass index (BMI) [17]. Indeed, OSA patients with ESRD tend to have a lower mean BMI than OSA patients with normal

* Corresponding author at: University Health Network Toronto General Hospital, 9N-943, 200 Elizabeth Street, Toronto, ON M5G 2C4, Canada. Tel.: +1 416 340 4719; fax: +1 416 340 4197.

E-mail address: douglas.bradley@utoronto.ca (T.D. Bradley).

renal function [2,17,18]. The severity of OSA has been shown to be directly related to the degree of overnight rostral fluid shift from the legs in both heart failure [19] and ESRD patients [10]. In ESRD patients, the volume of fluid accumulating in the legs during the day is likely to be greater than in the general population. Consequently, it is likely that fluid overload and increased overnight fluid shift from the legs play important roles in the pathogenesis of OSA in this population. As fluid overload and rostral fluid shift might distend the LA and increase fluid accumulation in the neck, we postulated that in ESRD patients, LA size would be related to the degree of overnight rostral fluid shift and that, in turn, OSA severity would be related to LA size. To test this hypothesis, we examined LA size determined by echocardiography in relation to OSA severity as determined by polysomnography (PSG) in patients with ESRD.

2. Methods

2.1. Subjects

Inclusion criteria were patients with ESRD at least 18 years of age undergoing thrice-weekly hemodialysis at the University Health Network Toronto General Hospital. Patients were recruited consecutively irrespective of symptoms of sleep apnea. Exclusion criteria were patients who were already treated for OSA or had an LV ejection fraction (LVEF) of <45%.

2.2. Protocol

2.2.1. Echocardiography

As part of routine clinical care in the hemodialysis unit, all patients with ESRD undergo echocardiography yearly. Transthoracic echocardiography was performed using a sector array between 1 and 5 Hz. Two-dimensional echocardiographic images were acquired from the parasternal long and short axes, apical long axis, apical four-chamber, apical two-chamber, and subcostal views.

Variables recorded were LVEF, LV mass, and LA size. LVEF was measured using the biplane Simpson's method using the apical four-chamber and apical two-chamber views. LA size was estimated by measuring the maximum internal anteroposterior diameter of the LA, using the parasternal long axis view. LV mass was estimated using the American Society of Echocardiography (ASE)-recommended formula: $LV\ mass = 0.8 \times (1.04 (LVDD + PWTD + IVSTD)^3 - (LVDD)^3) + 0.6\ g$, where LVDD is the LV internal diameter in diastole, PWTD is the posterior wall thickness in diastole, and IVSTD is the interventricular septum thickness in diastole. LA size index and LV mass index were calculated by expressing LA size and LV mass, respectively, per unit body surface area.

All echocardiograms were performed and reported prior to PSG so that the sonographer and reporting cardiologist were unaware of the patient's PSG findings and whether or not they had OSA.

2.2.2. Polysomnography

All subjects underwent overnight PSG the day before dialysis. Prior to PSG, demographic characteristics, medical history, and prescribed medications were recorded. PSG was performed with the use of standard techniques and scoring criteria for sleep stages and arousals from sleep [20]. All subjects slept on a single pillow with the bed flat. Thoracoabdominal motion was monitored by respiratory inductance plethysmography, and nasal airflow was monitored by nasal pressure cannulae (Binaps model 5500; Salter Labs, Arvin, CA, USA). Arterial oxyhemoglobin saturation (SaO₂) was monitored by oximetry. Obstructive apnea was defined as a >90% reduction of tidal volume for ≥10 s with thoracoabdominal motion, and obstructive hypopnea was defined as a 50–90% reduction in tidal volume from baseline for ≥10 s with out-of-phase thoracoabdominal motion or airflow limitation on nasal pressure. Apneas were clas-

sified as central in the absence of thoracoabdominal motion, and hypopneas were classified as central in the presence of in-phase thoracoabdominal motion and without airflow limitation on nasal pressure. Although these methods for classifying hypopneas as obstructive or central are in keeping with the criteria recommended by the American Academy of Sleep Medicine (AASM) for scoring hypopneas [21] and in most instances the distinction is clear, in others it is not. Thus, these techniques provide a reasonable approximation of the numbers of obstructive and central hypopneas. The apnea–hypopnea index (AHI) was calculated as the number of apneas and hypopneas per hour of sleep. Signals were recorded on a computerized sleep recording system (Sandman; Nellcor Puritan Bennett Ltd, Ottawa, Canada) and scored by personnel blind to measurements of leg fluid volume (LFV), neck circumference (NC), and echocardiographic findings.

2.2.3. LFV And neck circumference

A subset of patients agreed to have assessments of LFV and NC. With subjects instrumented for PSG, lying awake and supine with the legs straight, LFV was assessed by measuring the impedance to electrical flow between the electrodes placed on the ankle and upper thigh of the right leg by a bioelectrical impedance device (Xitron Hydra, model 4200, Xitron Technologies Inc., San Diego, CA, USA), as previously described [19,22–24]. This well-validated technique [25,26] uses the impedance to electric current within a body segment to measure its fluid content. Alterations in the fluid content of tissues cause proportional changes in impedance. NC was measured at the superior border of the cricothyroid cartilage with a tape measure. Lines drawn at this level ensured that measurements before and after sleep were made at the same place. On awakening the next morning, measurements made before sleep were repeated. The differences between LFV and NC, before and after sleep, were considered as the overnight changes in LFV (Δ LFV) and NC (Δ NC). Measurements of LFV and NC were made prior to scoring of the PSG by personnel unaware of the AHI.

The protocol was approved by the Research Ethics Board of the University Health Network and Toronto Rehabilitation Institute, and all subjects provided written informed consent before participation.

2.3. Statistical analysis

Relationships between single variables were examined by the Pearson correlation coefficient. Multivariable analysis was also undertaken with AHI as the dependent variable and with age, BMI, presence or absence of hypertension, LV mass, and LA size index as the independent variables using multiple stepwise linear regression with $p < 0.05$ to enter and $p > 0.1$ to remove. A similar multivariable analysis was undertaken with LA size index as the dependent variable and with Δ LFV, age, BMI, and presence or absence of hypertension as the independent variables. Data are presented as mean \pm standard deviation (SD) unless indicated otherwise. A p value <0.05 was considered significant. Analyses were performed with the use of SPSS 21.0.1 (SPSS Inc., Chicago, IL, USA).

3. Results

3.1. Characteristics of the patients

Forty ESRD patients underwent PSG and an echocardiogram was performed. Their characteristics are shown in Table 1. The study population was receiving adequate dialysis, as indicated by a percent reduction of urea >65% in all. The great majority of apneas and hypopneas were obstructive. A subset of the last 21 patients underwent LFV assessment.

Table 1
Characteristics of the patients who underwent polysomnography and echocardiography.

	n = 40
Men : Women	22:18
Age	47.1 ± 15.9
BMI, kg/m ²	27.5 ± 6.7
Reduction of urea, %	76.1 ± 10.2
AHI, events/h of sleep	25.1 ± 23.4
Obstructive AHI	21.7 ± 21.0
Central AHI	2.5 ± 4.6
LVEF%	64.4 ± 7.1
LV mass, g	204.1 ± 85.6
Left atrial size, mm	39.7 ± 7.8
Left atrial size index, mm/m ²	22.0 ± 4.9
Hypertension (n)	24
Atrial fibrillation (n)	1
Diabetes (n)	13

Abbreviations: BMI = body mass index, AHI = apnea–hypopnea index, LVEF = left ventricular ejection fraction.
Data are presented as mean ± standard deviation unless otherwise indicated.

3.2. Polysomnographic and echocardiographic data

Among the 40 patients, there was a significant correlation between AHI and LA size ($r = 0.626$, $p < 0.001$). In men, this relationship persisted ($r = 0.617$, $p = 0.002$) and became even stronger when LA size was indexed to body surface area ($r = 0.743$, $p < 0.001$, Fig. 1). In women, there was no significant relationship between AHI and LA size ($r = 0.434$, $p = 0.07$) or AHI and LA size index ($r = 0.325$, $p = 0.30$). In men, there was also a significant relationship between LV mass index and AHI ($r = 0.697$, $p = 0.001$) that was not seen in women ($r = 0.219$, $p = 0.38$). Stepwise multiple regression analysis in men revealed that the only factor that correlated independently with the AHI was the LA size index ($r = 0.781$, $r^2 = 0.609$, $p < 0.001$).

3.3. LfV measurements

Twenty-one patients underwent LFV assessments in addition to PSG and echocardiography. There were nine men and 12 women whose age was 43.8 ± 14.6 years, BMI was 27.1 ± 7.3 kg/m², and AHI




Fig. 2. Relationship between left atrial (LA) size index and overnight change in leg fluid volume (ΔLFV) in men.

was 18.3 ± 21.8 events/h of sleep. Among the men, there was a strong inverse correlation between LA size index and ΔLFV ($r = -0.739$, $p = 0.02$, Fig. 2) and also between AHI and ΔLFV ($r = -0.863$, $p = 0.003$, Fig. 3). Stepwise multiple regression analysis, in men, revealed that the only factor that correlated independently with the LA size index was ΔLFV ($r = -0.739$, $r^2 = 0.546$, $p = 0.02$). In women, there was a significant relationship between LA size and ΔLFV ($r = 0.601$, $p = 0.04$), but no relationship between LA size index and ΔLFV ($r = 0.077$, $p = 0.81$). There was also no relationship between AHI and ΔLFV in women ($r = 0.028$, $p = 0.93$). In addition, in men, there were significant relationships between AHI and LA size ($r = 0.719$, $p = 0.03$) and between AHI and LA size index ($r = 0.833$, $p = 0.005$). However, in women, these relationships were not significant ($r = 0.389$, $p = 0.21$ and $r = 0.325$, $p = 0.3$, respectively). There was no relationship between ΔLFV and LV mass index in men or women. In men, there was a direct relationship between ΔNC and LA size index ($r = 0.703$, $p = 0.04$) that was not seen in women ($r = 0.065$, $p = 0.84$).




Fig. 1. Relationship between apnea–hypopnea index (AHI) and left atrial (LA) size index in men.




Fig. 3. Relationship between apnea–hypopnea index (AHI) and change in overnight leg fluid volume (ΔLFV) in men.

4. Discussion

This study provides novel insights into potential mechanisms linking fluid retention and overnight rostral fluid shift with cardiac structure and OSA severity in ESRD patients on conventional hemodialysis. We found, in men, that there was a strong, direct relationship between severity of OSA, assessed by the AHI, and LA size. We also showed in men that there were strong relationships between the Δ LFV and both the LA size and the severity of OSA. Furthermore, in men, there was a relationship between the overnight change in NC and LA size. In women, however, there were no significant relationships between LA size and OSA severity, NC, or Δ LFV.

Fluid retention in ESRD is a major therapeutic challenge as it can lead to leg edema, pulmonary congestion, and systemic hypertension [27–29]. It is also associated with increased cardiovascular mortality, such that higher inter-dialytic weight gain is associated with increased cardiovascular deaths and worse survival [15]. However, the mechanisms by which fluid retention contributes to increased mortality have not been elucidated. Our study suggests that Δ LFV, LA size, and OSA severity may be involved in this relationship. In ESRD patients, it is possible that not only could fluid moving out of the legs overnight shift rostrally into the neck but it could also redistribute to the thorax and heart where it may distend the LA. This may, in turn, lead to an increase in central venous pressure, further fluid accumulation in the neck, an increase in pharyngeal collapsibility, and an increase in OSA severity. The observation that increased fluid removal via nocturnal peritoneal dialysis versus conventional continuous ambulatory dialysis leads to an increase in pharyngeal size and a reduction in AHI is in keeping with this proposal [30]. This possibility is further supported by our finding in this study of a strong relationship between LA size and Δ NC.

Increased LA size predicts increased cardiovascular mortality in ESRD patients, independently of LV mass or systolic dysfunction [5,31]. The LA can enlarge secondary to other cardiac pathology such as LV dysfunction, LV hypertrophy, or mitral valve disease [13,14]. However, its thin-walled structure makes it very susceptible to distension from pressure and volume overload, even in the absence of other cardiac pathology [13]. In this study, patients were excluded if their LVEF was <45% and their mean LVEF of 64 % is well within normal limits indicating normal systolic function. In addition, no patient had significant mitral or other valvular heart disease. There was a relationship between LA size and LV mass. However, while LA size index correlated strongly with Δ LFV, LV mass and LV mass index did not. The strong relationships seen between LA size and Δ LFV and between the AHI and LA size suggest that increased mortality associated with LA size may be at least partly related to the degree of overnight fluid shift from the legs and to the presence and severity of OSA.

It is also possible that diastolic dysfunction, which is common in patients with ESRD [32], contributed to the relationships among fluid overload, overnight fluid shift, LA size, and sleep apnea severity. However, diastolic dysfunction, itself, is profoundly influenced by alterations in preload and as this is constantly changing in ESRD because of pronounced fluid volume shifts from day to day due to dialysis, it is very difficult to reliably diagnose diastolic dysfunction in ESRD patients by echocardiography alone. Furthermore, in the setting of preserved LV systolic function, recent guidelines advise that flow Doppler measurements alone are not adequate to identify diastolic dysfunction and recommend the use of tissue Doppler at the very least [33]. However, tissue Doppler was not performed in this study. In any case, given the limitations of echocardiography in distinguishing true diastolic dysfunction from volume overload, it is likely that more invasive investigations would be required to make a definitive diagnosis of diastolic dysfunction in ESRD patients such as ours.

OSA is much more common in men than women in the general population as well as in ESRD patients [8,34]. Various factors, such as a shorter pharynx, different neck fat distribution, genioglossus activity, and hormonal status, may reduce women's susceptibility to OSA [35–37]. None, however, can fully explain the difference in OSA prevalence between men and women [37]. While we found a direct relationship between Δ LFV and AHI in men, we found no such relationship in women, consistent with our previous findings in patients with heart failure [38]. Previous studies have also shown that for a similar degree of fluid shift from the legs, the increase in NC and UA collapsibility was much greater in men than in women [19,39]. In women, a greater proportion of the fluid shifting from the legs may be redistributed to areas aside from the neck such as the large venous plexi around the uterus and ovaries. It is therefore possible that differences in the pattern of overnight fluid redistribution might be another factor that contributes to the lower prevalence of OSA in women than in men.

In summary, we report novel data demonstrating that LA size is related to both Δ LFV and severity of OSA in men but not in women with ESRD. Thus, in men, the relationship between LA size and mortality in ESRD may also be linked to degrees of fluid overload, overnight rostral fluid shift, and severity of OSA. Further studies of overnight fluid redistribution that include measurements of fluid volumes of the abdomen, thorax, and neck may help determine if the difference in OSA prevalence between men and women can be explained by sex-specific differences in fluid redistribution patterns. Finally, given the relationships among overnight fluid shift from the legs and both LA size and OSA severity, and their potential contribution to increased mortality in this high-cardiovascular risk group, our results highlight the importance of determining whether removal of excess fluid by intensified dialysis or ultrafiltration would reduce LA size and OSA severity in ESRD patients.

Conflict of interest

The ICMJE Uniform Disclosure Form for Potential Conflicts of Interest associated with this article can be viewed by clicking on the following link: <http://dx.doi.org/10.1016/j.sleep.2014.07.001>.

Acknowledgments

This work was supported by operating grant MOP-82731 from the Canadian Institutes of Health Research. OD Lyons is supported by a joint Canadian Thoracic Society/European Respiratory Society Peter Macklem Research Fellowship, CT Chan by the R. Fraser Elliot Chair in Home Dialysis, RM Elias by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Brazil, Toronto Rehabilitation Institute and Toronto General Hospital, and TD Bradley by the Clifford Nardal Chair in Sleep Apnea and Rehabilitation Research.

References

- [1] London GM. Cardiovascular disease in chronic renal failure: pathophysiologic aspects. *Semin Dial* 2003;16(2):85–94. PubMed PMID: 12641870.
- [2] Zoccali C, Mallamaci F, Tripepi G. Nocturnal hypoxemia predicts incident cardiovascular complications in dialysis patients. *J Am Soc Nephrol* 2002;13(3):729–33. PubMed PMID: 11856778.
- [3] Zoccali C, Mallamaci F, Tripepi G. Sleep apnea in renal patients. *J Am Soc Nephrol* 2001;12(12):2854–9. PubMed PMID: 11729258.
- [4] Benjamin EJ, D'Agostino RB, Belanger AJ, Wolf PA, Levy D. Left atrial size and the risk of stroke and death. The Framingham Heart Study. *Circulation* 1995;92(4):835–41. PubMed PMID: 7641364.
- [5] Shizuku J, Yamashita T, Ohba T, Kabaya T, Nitta K. Left atrial volume is an independent predictor of all-cause mortality in chronic hemodialysis patients. *Intern Med* 2012;51(12):1479–85. PubMed PMID: 22728478.
- [6] Hanly PJ, Pierratos A. Improvement of sleep apnea in patients with chronic renal failure who undergo nocturnal hemodialysis. *N Engl J Med* 2001;344(2):102–7. PubMed PMID: 11150360.
- [7] Kimmel PL, Miller G, Mendelson WB. Sleep apnea syndrome in chronic renal disease. *Am J Med* 1989;86(3):308–14. PubMed PMID: 2919612.

- [8] Argekar P, Griffin V, Litaker D, Rahman M. Sleep apnea in hemodialysis patients: risk factors and effect on survival. *Hemodial Int* 2007;11(4):435–41. PubMed PMID: 17922741.
- [9] Tang SC, Lam B, Yao TJ, Leung WS, Chu CM, Ho YW, et al. Sleep apnea is a novel risk predictor of cardiovascular morbidity and death in patients receiving peritoneal dialysis. *Kidney Int* 2010;77(11):1031–8. PubMed PMID: 20237456.
- [10] Elias RM, Bradley TD, Kasai T, Motwani SS, Chan CT. Rostral overnight fluid shift in end-stage renal disease: relationship with obstructive sleep apnea. *Nephrol Dial Transplant* 2012;27(4):1569–73. PubMed PMID: 22058175.
- [11] Bradley TD, Floras JS. Obstructive sleep apnoea and its cardiovascular consequences. *Lancet* 2009;373(9657):82–93. PubMed PMID: 19101028.
- [12] Rosca M, Lancellotti P, Popescu BA, Pierard LA. Left atrial function: pathophysiology, echocardiographic assessment, and clinical applications. *Heart* 2011;97(23):1982–9. PubMed PMID: 22058287.
- [13] Leung DY, Boyd A, Ng AA, Chi C, Thomas L. Echocardiographic evaluation of left atrial size and function: current understanding, pathophysiologic correlates, and prognostic implications. *Am Heart J* 2008;156(6):1056–64. PubMed PMID: 19032999.
- [14] Abhayaratna WP, Seward JB, Appleton CP, Douglas PS, Oh JK, Tajik AJ, et al. Left atrial size: physiologic determinants and clinical applications. *J Am Coll Cardiol* 2006;47(12):2357–63. PubMed PMID: 16781359.
- [15] Kalantar-Zadeh K, Regidor DL, Kovesdy CP, Van Wyck D, Bunnapradist S, Horwich TB, et al. Fluid retention is associated with cardiovascular mortality in patients undergoing long-term hemodialysis. *Circulation* 2009;119(5):671–9. PubMed PMID: 19171851. Pubmed Central PMCID: 2773290.
- [16] Yumino D, Wang H, Floras JS, Newton GE, Mak S, Ruttanaumpawan P, et al. Prevalence and physiological predictors of sleep apnea in patients with heart failure and systolic dysfunction. *J Card Fail* 2009;15(4):279–85. PubMed PMID: 19398074.
- [17] Tada T, Kusano KF, Ogawa A, Iwasaki J, Sakuragi S, Kusano I, et al. The predictors of central and obstructive sleep apnoea in haemodialysis patients. *Nephrol Dial Transplant* 2007;22(4):1190–7. PubMed PMID: 17277346.
- [18] Beecroft JM, Pierratos A, Hanly PJ. Clinical presentation of obstructive sleep apnea in patients with end-stage renal disease. *J Clin Sleep Med* 2009;5(2):115–21. PubMed PMID: 19968043. Pubmed Central PMCID: 2670329.
- [19] Yumino D, Redolfi S, Ruttanaumpawan P, Su MC, Smith S, Newton GE, et al. Nocturnal rostral fluid shift: a unifying concept for the pathogenesis of obstructive and central sleep apnea in men with heart failure. *Circulation* 2010;121(14):1598–605. PubMed PMID: 20351237.
- [20] Rechtschaffen A, Kales A. A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects. Los Angeles, CA: UCLA Brain Information Service/Brain Research Institute; 1968.
- [21] Berry RB, Budhiraja R, Gottlieb DJ, Gozal D, Iber C, Kapur VK, et al. Rules for scoring respiratory events in sleep: update of the 2007 AASM manual for the scoring of sleep and associated events. Deliberations of the sleep apnea definitions task force of the American academy of sleep medicine. *J Clin Sleep Med* 2012;8(5):597–619. PubMed PMID: 23066376. Pubmed Central PMCID: 3459210.
- [22] Redolfi S, Yumino D, Ruttanaumpawan P, Yau B, Su MC, Lam J, et al. Relationship between overnight rostral fluid shift and Obstructive Sleep Apnea in nonobese men. *Am J Respir Crit Care Med* 2009;179(3):241–6. PubMed PMID: 19011149.
- [23] Chiu KL, Ryan CM, Shiota S, Ruttanaumpawan P, Arzt M, Haight JS, et al. Fluid shift by lower body positive pressure increases pharyngeal resistance in healthy subjects. *Am J Respir Crit Care Med* 2006;174(12):1378–83. PubMed PMID: 16998093.
- [24] Shiota S, Ryan CM, Chiu KL, Ruttanaumpawan P, Haight J, Arzt M, et al. Alterations in upper airway cross-sectional area in response to lower body positive pressure in healthy subjects. *Thorax* 2007;62(10):868–72. PubMed PMID: 17442706. Pubmed Central PMCID: 2094267.
- [25] Demura S, Sato S, Kitabayashi T. Percentage of total body fat as estimated by three automatic bioelectrical impedance analyzers. *J Physiol Anthropol Appl Human Sci* 2004;23(3):93–9. PubMed PMID: 15187381.
- [26] Zhu F, Kuhlmann MK, Kotanko P, Seibert E, Leonard EF, Levin NW. A method for the estimation of hydration state during hemodialysis using a calf bioimpedance technique. *Physiol Meas* 2008;29(6):S503–16. PubMed PMID: 18544816.
- [27] Katzarski KS. Monitoring of blood volume during haemodialysis treatment of acute renal and multiple organ failures. *Nephrol Dial Transplant* 1996;11(Suppl. 8):20–3. PubMed PMID: 9044336.
- [28] Charra B, Laurent G, Chazot C, Calemard E, Terrat JC, Vanel T, et al. Clinical assessment of dry weight. *Nephrol Dial Transplant* 1996;11(Suppl. 2):16–19. PubMed PMID: 8803988.
- [29] Kraemer M, Rode C, Wizemann V. Detection limit of methods to assess fluid status changes in dialysis patients. *Kidney Int* 2006;69(9):1609–20. PubMed PMID: 16501488.
- [30] Tang SC, Lam B, Lai AS, Pang CB, Tso WK, Khong PL, et al. Improvement in sleep apnea during nocturnal peritoneal dialysis is associated with reduced airway congestion and better uremic clearance. *Clin J Am Soc Nephrol* 2009;4(2):410–18. PubMed PMID: 19118118. Pubmed Central PMCID: 2637593.
- [31] Tripepi G, Benedetto FA, Mallamaci F, Tripepi R, Malatino L, Zoccali C. Left atrial volume monitoring and cardiovascular risk in patients with end-stage renal disease: a prospective cohort study. *J Am Soc Nephrol* 2007;18(4):1316–22. PubMed PMID: 17360950.
- [32] de Bie MK, Ajmone Marsan N, Gaasbeek A, Bax JJ, Groeneveld M, Gabreels BA, et al. Left ventricular diastolic dysfunction in dialysis patients assessed by novel speckle tracking strain rate analysis: prevalence and determinants. *Int J Nephrol* 2012;2012:963504. PubMed PMID: 22649726. Pubmed Central PMCID: 3357524.
- [33] Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, et al. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *J Am Soc Echocardiogr* 2009;22(2):107–33. PubMed PMID: 19187853.
- [34] Young T, Shahar E, Nieto FJ, Redline S, Newman AB, Gottlieb DJ, et al. Predictors of sleep-disordered breathing in community-dwelling adults: the Sleep Heart Health Study. *Arch Intern Med* 2002;162(8):893–900. PubMed PMID: 11966340.
- [35] Malhotra A, Huang Y, Fogel RB, Pillar G, Edwards JK, Kikinis R, et al. The male predisposition to pharyngeal collapse: importance of airway length. *Am J Respir Crit Care Med* 2002;166(10):1388–95. PubMed PMID: 12421747.
- [36] Chin CH, Kirkness JP, Patil SP, McGinley BM, Smith PL, Schwartz AR, et al. Compensatory responses to upper airway obstruction in obese apneic men and women. *J Appl Physiol* 2012;112(3):403–10. PubMed PMID: 22096117. Pubmed Central PMCID: 3289429.
- [37] Lin CM, Davidson TM, Ancoli-Israel S. Gender differences in obstructive sleep apnea and treatment implications. *Sleep Med Rev* 2008;12(6):481–96. PubMed PMID: 18951050. Pubmed Central PMCID: 2642982.
- [38] Kasai T, Motwani SS, Yumino D, Mak S, Newton GE, Bradley TD. Differing relationship of nocturnal fluid shifts to sleep apnea in men and women with heart failure. *Circ Heart Fail* 2012;5(4):467–74. PubMed PMID: 22679060.
- [39] Su MC, Chiu KL, Ruttanaumpawan P, Shiota S, Yumino D, Redolfi S, et al. Difference in upper airway collapsibility during wakefulness between men and women in response to lower-body positive pressure. *Clin Sci* 2009;116(9):713–20. PubMed PMID: 19007335.